CHAPTER FIVE

Oocyte Meiotic Spindle Assembly and Function

Aaron F. Severson[*], George von Dassow[†], Bruce Bowerman[{,1]
[*]Department of Biological, Geological, and Environmental Sciences, Center for Gene Regulation in Health and Disease, Cleveland State University, Cleveland, Ohio, USA
[†]Oregon Institute of Marine Biology, University of Oregon, Charleston, Oregon, USA
[{]Institute of Molecular Biology, University of Oregon, Eugene, Oregon, USA
[1]Corresponding author: e-mail address: bowerman@uoregon.edu

Contents

1. Introduction 66
2. Maintain Parity or Pare Down: Mitotic Versus Meiotic Cell Division 66
3. Centrosomes: The Stars of Mitotic Spindle Poles 70
4. The Curious Structure of (Some) Oocyte Meiotic Spindles: Acentriolar Poles 71
5. The Curious Structure of Oocyte Meiotic Spindles, Part 2: Tiled Microtubules 75
6. Acentriolar Oocyte Meiotic Spindle Assembly: Microtubule Origins 77
7. Acentriolar Oocyte Meiotic Spindle Assembly, Part 2: Pole Composition 79
8. Acentriolar Oocyte Meiotic Spindle Assembly, Part 3: Pole Assembly 81
9. Kinetochore Function and Pole Coalescence 84
10. A Pushy View of Kinetochore Function and Chromosome Dynamics During Oocyte Meiosis 87
11. An Alternative Model for Chromosome Movement in *C. elegans* Oocytes: Sides Matter 88
12. Advancing Our Understanding of Oocyte Meiotic Spindle Assembly: Genetics and Live Cell Imaging 91

Acknowledgments 92
References 92

Abstract

Gametogenesis in animal oocytes reduces the diploid genome content of germline precursors to a haploid state in gametes by discarding half of the duplicated chromosomes through a sequence of two meiotic cell divisions called meiosis I and II. The assembly of the microtubule-based spindle structure that mediates this reduction in genome content remains poorly understood compared to our knowledge of mitotic spindle assembly and function. In this review, we consider the diversity of oocyte meiotic spindle assembly and structure across animal phylogeny, review recent advances in our understanding of how animal oocytes assemble spindles in the absence of the centriole-based microtubule-organizing centers that dominate mitotic spindle assembly, and

Current Topics in Developmental Biology, Volume 116
ISSN 0070-2153
http://dx.doi.org/10.1016/bs.ctdb.2015.11.031

2016 Elsevier Inc.
All rights reserved.


Sister chromatid cohesion
Centromere/kinetochore

and are formed by the polymerization of and -tubulin dimers (

destroyed. While meiotic and mitotic cohesins share several subunits, certain

the PCM of each centrosome, the microtubules they nucleate project outward in all directions, forming large and radially symmetric asters. Ultimately, some microtubules from each centrosome capture and align chromosomes at the metaphase plate, and then mediate chromosome segregation to the poles during anaphase.

Importantly, centrosome duplication is tightly regulated during the cell cycle (Firat-Karalar & Stearns, 2014). After mitosis, each daughter cell receives one pair of centrioles that separate. Each centriole then produces a new, orthogonally oriented daughter centriole, restoring the seeds of bipolarity for the next mitotic cell division (Gönczy, 2012). Intriguingly, centrioles also serve as the basal bodies from which cilia and flagella grow at the surface of interphase cells. These roles—as centrosome organizers or basal bodies—appear to be mutually exclusive in all animal cells.

4. THE CURIOUS STRUCTURE OF (SOME) OOCYTE MEIOTIC SPINDLES: ACENTRIOLAR POLES

In contrast to the spindles formed during mitosis, oocyte meiotic spindles in many animals—including humans and the laboratory models of mice, frogs, fruit flies, and nematodes—entirely lack centrioles (Dumont & Desai, 2012; Howe & FitzHarris, 2013; Müller-Reichert et al., 2010; Ohkura, 2015). Nevertheless, bipolar spindles still assemble in the absence of these key organizers of the mitotic spindle. Moreover, even oocytes that retain centrioles at meiotic spindle poles typically lose them during or shortly after meiosis, and thus most animal eggs ultimately lack functional centrioles.

But why eliminate centrioles before meiosis? Based on the limited comparative data available, it seems likely that primitive animal oocytes retained centrioles and centrosome-based meiotic spindle pole function, complete with astral microtubules. For example, during meiosis I in sea star and sea urchin (echinoderm) oocytes, each spindle pole harbors a pair of centrioles, as in mitosis, but centriole duplication does not occur after meiosis I. Each pole in meiosis II therefore has only a single centriole (Nakashima & Kato, 2001; Sluder et al., 1989). Experimental analysis of sea star oocytes indicates that the centrioles inherited by polar bodies are fully functional and replicable, whereas the single centriole that remains in the egg after meiosis II is apparently nonreplicable, and presumably eventually degenerates (Sluder et al., 1989; Uetake et al., 2007). The two pairs of centrioles derived from the sperm then mediate the first mitotic division of the zygote. Thus, in sea stars the oocyte's complement of replication-competent centrioles is discarded into polar bodies. Indeed, the failure of polar body emission enables parthenogenetic development due to the retention of functional, reproductive centrosomes (Washitani-Nemoto et al., 1994).

Echinoderms are deuterostomes, like vertebrates, and another echinoderm group, the sea cucumbers, also have centriolar spindles (Hollers, 1981). However, chordates, including the tunicates, all appear to assemble acentriolar spindles during oocyte meiosis (Sawada & Schatten, 1988). Which mode is primitive for animals?

Many invertebrate protostomes also retain centrioles through oocyte meiosis. Centrioles have either been detected using transmission electron microscopy (TEM) or have been inferred to exist because of the presence of extensive astral microtubule arrays at meiotic spindle poles. Therefore, centriolar oocyte spindles may be the ancestral condition. In the clam *Spisula solidissima*, the fertilized oocyte contains three centrosomes during prometaphase of meiosis I—one from the sperm and two from the oocyte (Wu & Palazzo, 1999). As in echinoderms, one oocyte centrosome with paired centrioles is extruded into the first polar body at the end of meiosis I. A second oocyte centrosome, with an unpaired centriole, is extruded into the second polar body at the end of meiosis II. Thus, the zygote inherits one maternal centrosome with a single, unduplicated centriole and one paternal centrosome with a duplicated centriole. While the fate of the remaining oocyte centrosome is unknown, subsequent mitotic divisions use the sperm-derived centrioles. Remarkably, the sperm centrosome appears to be repressed during oocyte meiosis: both α-tubulin and γ-tubulin become undetectable by metaphase of meiosis I. Similar suppression of the sperm

centrosome is observed in annelids and echinoderms (Stephano & Gould, 2000). Spisula likely typifies the entire Lophotrochozoan supergroup; although TEM has been applied to identify centrioles in only a few species in this group, the presence of large asters around meiotic spindle poles in oocytes of numerous Lophotrochozoan species implies their presence (Crowder et al., 2015b). Meanwhile, although insect and C. elegans oocytes feature well-studied acentriolar (and largely anastral) meiotic spindles, this is unlikely to be primitive for the Ecdysozoan supergroup as the oocyte meiotic spindles of some crustaceans have large astral, centriolar poles (Goudeau & Goudeau, 1986; Goudeau & Lachaise, 1980; Lindsay et al., 1992).

Although comparative data are limited, acentriolar spindle assembly pathways likely evolved multiple times in animals, and nearly exclusively for use in oocyte meiosis. This in turn suggests that similar selective pressures may have favored the transition to acentriolar oocyte spindles in these lineages. One adaptive advantage might have been to obviate the book-keeping involved in sorting centrioles from different sources into polar bodies versus eggs. Although this seems like a minor mechanistic burden, any distinctions that enable predictable centriole sorting could in principle be interpreted by other agents, making meiosis more susceptible to cheating (i.e., meiotic drive).

Another often cited rationale for acentriolar meiosis is that small and largely anastral meiotic spindles might minimize the size of polar bodies, the discard products of meiosis. The extreme asymmetry of the two divisions that occur during oocyte meiosis ensures that the zygote inherits almost all of the maternally synthesized gene products, biosynthetic machinery, and fuel that sustain early embryonic development. For example, in C. elegans oocytes, very small meiotic spindles assemble in close proximity to the cortex (Fig. 2). While the zygote itself is approximately 50 μm, the oocyte meiotic spindles initially are roughly 8 μm in length and by meta-phase shorten to only 5 μm (Albertson & Thomson, 1993; McNally & McNally, 2005; McNally et al., 2006; Yang et al., 2003). By contrast, the first mitotic spindle is much larger (Fig. 2), with astral microtubules spanning the entire length and width of the one-cell zygote (Müller-Reichert et al., 2010). Only a small number of short astral microtubules have been detected during oocyte meiosis in C. elegans; these microtubules may be important for the microtubule motor-dependent translocation and rotation of the spindle prior to anaphase (Crowder et al., 2015a; Ellefson & McNally, 2009, 2011; Yang et al., 2005). Indeed, spindle positioning in C. elegans oocytes is micro-tubule-dependent but actin-independent (Yang et al., 2003). In contrast,

rotation of the acentriolar and likely asterless oocyte spindles in mouse oocytes requires actin and myosin but appears to be independent of any microtubule motors (Maro et al., 1984; Wang et al., 2011). It is not clear whether acentriolar oocyte meiotic spindles in Drosophila have any astral microtubules (Skold et al., 2005); however, asters are detectable in Xenopus oocytes (Gard, 1992), although it is not known if they are involved in spindle positioning.

While small not

enough cytoplasm, they can continue mitosis (Saiki & Hamaguchi, 1997). If such cells were incorporated into the embryo it would potentially violate these imperatives. The chance of this happening is remote in free-spawning invertebrates, but is perhaps higher in encapsulated, brooded, or placental embryos. Depriving polar bodies of centrioles might forestall this possibility. That this is a plausible risk is shown by the rare instances, in humans, of polar body fertilization, giving rise to a chimera or even a twin (Bieber et al., 1981).

Finally, consider the unusual way the freshwater clam *Corbicula leana*, a hermaphroditic triploid, begins a new life. Centrosomes with typical centriole pairs mediate assembly of the oocyte meiosis I spindle, but the spindle remains parallel to the plasma membrane and both poles and all maternal chromosomes are extruded simultaneously into a pair of polar bodies (Komaru et al., 2000). Thus, development in this species is androgenetic: the zygote chromosomes come entirely from the sperm. This is doubtless among the most extreme variants of meiosis, but underscores the extent to which this biological process, so central to all eukaryotic sexuality, evolves in concert with animal life histories.

The intriguing variation of meiotic mechanisms across animal phylogeny seems largely neglected, despite its obvious possible adaptive significance. Nematodes provide one avenue for exploring this remarkable example of evolutionary cell biology. Their oocytes are amenable to live imaging, and many different species have been isolated, often with fully sequenced genomes (Frauchle et al., 2009; Farhadifar et al., 2015; Phillips & Bowerman, 2015). Moreover, there is substantial cryptic genetic variation that impacts embryonic viability among wild *C. elegans* populations (Paaby et al., 2015). Perhaps, some of this genotypic variation influences oocyte spindle assembly and might shed light on the phenotypic variation observed at the family and phyla levels.

5. THE CURIOUS STRUCTURE OF OOCYTE MEIOTIC SPINDLES, PART 2: TILED MICROTUBULES

Another remarkable difference between mitotic and at least some oocyte meiotic spindles is the continuous versus discontinuous nature of the microtubules that span the distance between the poles and the spindle midzone and chromosomes (Fig. 2B). Although mitotic spindles may include some short tiled microtubules (Goshima & Kimura, 2010), long microtubules typically extend from each mitotic centrosome to the midzone

and beyond (White & Glotzer, 2012). In some cases, the long microtubules form crosslinked, antiparallel bundles that stabilize the mitotic spindle and ultimately constitute the central spindle and midbody remnant, structures that promote the initiation and completion of cytokinesis, respectively (

understand where these factors act. While a gradient of active Ran GTPase

2015; Ellefson & McNally, 2011; Gomes et al., 2013; McNally et al., 2006, 2014; O'Rourke et al., 2007; van der Voet et al., 2009; Wignall & Villeneuve, 2009). Thus far, -tubulin has not been clearly detected at poles but appears to be present diffusely throughout the oocyte spindle (McNally et al., 2006). While less is known about pole composition in Drosophila, augmin subunits and the conserved microtubule-associated proteins (MAPs) minispindles and D-TACC have been detected at mature poles (Colombie et al., 2013; Cullen & Ohkura, 2001). Drosophila -tubulin appears to be somewhat enriched at oocyte spindle poles but also colocalizes with microtubules throughout the prometaphase meiosis I spindle (Endow & Hallen, 2011; Hughes et al., 2011).

8. ACENTRIOLAR OOCYTE MEIOTIC SPINDLE ASSEMBLY, PART 3: POLE ASSEMBLY

A landmark advance in our understanding of oocyte meiotic spindle dynamics came from live cell imaging of pole assembly in mouse oocytes, which showed that multiple small pole foci coalesce to form a bipolar spindle structure (Schuh & Ellenberg, 2007). While earlier studies of fixed oocytes first documented this phenomenon (Carabatsos et al., 2000; Gueth-Hallonet et al., 1993; Palacios et al., 1993), Schuh and Ellenberg used live cell imaging to show that many small MTOCs are initially dispersed throughout the area surrounding the mouse oocyte chromosomes early in meiosis I. Over time these small foci coalesce to form a bipolar structure.

More recently, live imaging of Cep192 fusion to GFP has shown that early pole coalescence in mouse oocytes involves an early dispersal of fewer and larger MTOC foci into more and smaller foci, followed by a still mysterious coalescence (Clift & Schuh, 2015). An early phase of dispersal, prior to nuclear envelope breakdown, requires the polo kinase PLK1 and the minus-end directed microtubule motor dynein, in a kinae1i323(fu1Stchted)-1

more recent study using Cep192 as a marker for MTOC pole foci failed to detect a ball-like structure with MTOCs surrounded by chromosomes; rather the MTOCs and chromosomes were comingled (Clift & Schuh, 2015). In both studies, the subsequent dispersal of MTOCs was shown to require the kinesin-5 family member KIF11 (Fig. 3A) (Clift & Schuh, 2015; Schuh & Ellenberg, 2007). How the multiple, dispersed MTOCs coalesce to form two poles remains unknown.

In Xenopus extracts, discrete, small microtubule foci have not been reported as pole intermediates during the assembly of oocyte meiotic spindles. However, acentrosomal bipolar spindle organization does require a kinesin-5 family member (Eg5), as in mouse. Pole assembly also is thought to involve crosslinking of microtubule minus ends mediated by dynein and the coiled-coil protein NUMA (Heald et al., 1996; Merdes et al., 2000; Mitchison et al., 2005).

In C. elegans, live-imaging studies indicate that chromosomes are initially dispersed throughout a mass of microtubules, and the small ASPM-1 pole foci formed early in meiosis coalesce to form a bipolar structure (Fig. 3B) (Connolly et al., 2015). Unlike in mice and frogs, kinesin-5 (C. elegans BMK-1) is not required for bipolar spindle assembly during meiosis or mito-

assembly. However, *subito* mutants display transient instability of spindle poles, suggesting that Subito-mediated central spindle integrity may be important for promoting spindle bipolarity (Colombie et al., 2013; Jang et al., 2007).

In sum, the molecular pathways that promote pole coalescence and spin-

B Anaphase pushing model



Prometaphase Metaphase Anaphase

Spindle poles Ring proteins (CLASP) Kinetochore proteins (KNL-1)

C Lateral attachments model

Prometaphase Metaphase

during oocyte meiosis might lead to abnormal tension within the assembling spindle, and this imbalance in forces has been proposed to interfere with the coalescence of ASPM-1 foci into a bipolar structure (Connolly et al., 2015). Consistent with such a model, partial knockdown of components of the Ndc80 complex that mediates microtubule–kinetochore attachment rescues spindle bipolarity in klp-7( ) mutants.

While the importance of microtubule–kinetochore attachment for oocyte spindle assembly in other species remains largely unknown, roughly 90% of the chromosomes in mouse oocytes transiently form improper syntelic or merotelic attachments (Kitajima et al., 2011). Moreover, knockdown of the Ndc80 complex in both mouse and nematode oocytes results in spindle assembly defects (Breuer et al., 2010; Sun et al., 2010, 2011), suggesting that a role for kinetochores is conserved. Finally, expression of a dominant negative allele of Drosophila KLP10A, a kinesin-13/MCAK family member, results in disorganized or extra oocyte spindles poles (Zou et al., 2008), suggesting that a role for kinesin-13/MCAK in limiting pole number also may be conserved. It will be interesting to test whether MCAK-mediated removal of improper microtubule–kinetochore attachments is important for oocyte spindle pole coalescence in other organisms.

Figure 4—Cont'd depolymerase KLP-7/MCAK. Failure to disrupt these incorrect attachments in mutants with reduced KLP-7 function results in an imbalance in spindle tension that interferes with the coalescence of early pole foci and thus results in the assembly of multipolar spindles. (B) Microtubule–kinetochore interactions orient and align bivalents on the metaphase spindle, but at the metaphase-to-anaphase transition, the polar microtubule arrays disassemble, and a new population of microtubules forms between homologs and pushes them apart independently of kinetochores. The assembly of these microtubules is mediated by CLASP and other factors present in ring-shaped structures between each homolog pair at metaphase. When homologs separate, the rings are left behind at the metaphase plate. (C) Lateral interactions between microtubules and microtubule motor proteins mediate both chromosome congression to the metaphase plate and anaphase segregation to the poles. The chromokinesin KLP-19, localized to midbivalent rings, interacts with spindle microtubules to produce a polar ejection force that aligns bivalents at the metaphase plate. Subsequently, during anaphase, polar microtubules interact with dynein, a minus-end directed microtubule

10. A PUSHY VIEW OF KINETOCHORE FUNCTION AND CHROMOSOME DYNAMICS DURING OOCYTE MEIOSIS

The influence of microtubule–kinetochore attachment on oocyte spindle pole coalescence in *C. elegans* brings us to an intriguing controversy regarding the importance of kinetochores during oocyte meiosis (Fig. 4B and C). An important early observation was that depleting either components of the Ndc80 complex, responsible for the microtubule attachment activity of the kinetochore, or core kinetochore components such as KNL-1, had remarkably minor effects on meiotic spindle assembly and chromosome organization (Dumont et al., 2010). In live-imaging studies of microtubule and chromosome dynamics, using fusions of GFP and mCherry to -tubulin and a histone, respectively, kinetochore disruption was shown to cause only minor perturbations in spindle morphology, chromosome congression to the metaphase plate, and anaphase chromosome segregation to the poles. During meiosis I in wild-type *C. elegans* oocytes, spindles shorten substantially prior to anaphase, and during anaphase the poles rapidly disassemble and most spindle microtubules are detected between the separating chromosomes (Fig. 4B). Moreover, knockdown of factors required for the assembly or stability of these anaphase microtubules, such as the microtubule-stabilizing protein CLS-2/CLASP, resulted in both a substantial loss of the interchromosomal microtubules and severe defects in anaphase chromosome movements (Dumont et al., 2010). These findings led Dumont and Desai to propose a model in which microtubule polymerization between the segregating chromosomes pushes the chromosomes apart during anaphase of meiosis I and II, with little if any requirement for microtubule–kinetochore interactions and poleward pulling forces during anaphase (Fig. 4B).

Whether the apparent lack of a substantial role for kinetochores during anaphase in *C. elegans* oocytes is relevant to other species is not known. However, TEM analysis of fixed mouse oocytes during meiosis I and II clearly indicate that microtubules attach to kinetochores, and the kinetochore regions lead the anaphase movements of meiotic chromosomes toward the poles (Brunet et al., 1999). Microtubule–kinetochore attachments have also been observed in human oocytes, although the correction of improper syntelic and merotelic attachments appears to be remarkably inefficient (Holubcova et al., 2015). Furthermore, knockdowns of Ndc80

complex components in mouse oocytes cause substantial defects in spindle organization and chromosome segregation (Sun et al., 2010, 2011), and perturbations of mouse meiotic spindle assembly lead to activation of the kinetochore-based spindle assembly checkpoint (Marat et al., 2010; McGuinness et al., 2009). Nevertheless, very few microtubule–kinetochore attachments form prior to metaphase of meiosis I in mouse oocytes and microtubule–kinetochore attachments may not be required for chromosome congression to the metaphase plate (Brunet et al., 1999). Although mutational inactivation of -tubulin disrupts microtubule–kinetochore attachments in Drosophila oocytes, the role of these attachments in chromosome segregation is unknown (Hughes et al., 2011). Systematic investigations of how microtubule–kinetochore attachments, and kinetochore function more generally, influence spindle assembly and chromosome movement are needed to fully assess and compare the role of these structures during oocyte meiotic cell division in different animal species.

11. AN ALTERNATIVE MODEL FOR CHROMOSOME MOVEMENTS IN C. ELEGANS OOCYTES: SIDES MATTER

While kinetochore-independent microtubule polymerization has been proposed to push chromosomes apart during meiotic anaphase in C. elegans oocytes (Dumont et al., 2010), two other studies have suggested a very different model for chromosome congression and segregation (Muscat et al., 2015; Wignall & Villeneuve, 2009). In this model (Fig. 4C), kinetochores are proposed to have little if any role during either congression or segregation. Instead, lateral attachments of polar microtubules to chromosomes, and microtubule motor-mediated pushing and pulling forces, were proposed to move chromosomes during both congression to the metaphase plate and anaphase poleward segregation, respectively.

The first of these two studies demonstrated that the chromokinesin KLP-19 localizes to rings that encircle the midregions of each bivalent, that this midbivalent accumulation of KLP-19 requires the CPC, and that KLP-19 is important for chromosome congression to the metaphase plate (Wignall & Villeneuve, 2009). Knockdown of KLP-19 had relatively modest effects on chromosome orientation relative to the spindle poles, but congression of chromosomes to the metaphase plate was often defective. Moreover, when KLP-19 was knocked down in mutants that make monopolar spindles, chromosomes remained clustered around the monopole. In contrast, chromosomes were usually found close to microtubule ends, far from the

monopole, in oocytes with wild-type levels of KLP-19. Wignall and Villeneuve therefore proposed that lateral associations of spindle microtubules with meiotic chromosomes, together with KLP-19 motor activity, creates a polar ejection force that moves chromosomes toward the spindle midzone. Upon reaching the midzone, chromosomes encounter overlapping, antiparallel microtubules, and a balance of forces result in no net movement toward either pole, aligning chromosomes at the metaphase plate. Polar ejection forces have also been proposed to move chromosomes toward the metaphase plate in *Drosophila* oocytes, in the context of augmin-mediated microtubule polymerization, although whether these forces involve lateral microtubule attachments or end-on attachment at kinetochores was not addrest

the opposite effect—lagging chromosomes were observed during anaphase in bipolar spindles and chromosomes failed to move centripetally in monopolar spindles. However, it is important to note that KLP-19 accumulates on both the lateral and poleward faces of meiotic bivalents and on meiotic spindle microtubules (Powers et al., 2004). Similarly, dynein is broadly distributed throughout the meiotic spindle apparatus (Crowder et al., 2015a; Ellefson & McNally, 2009, 2011; van der Voet et al., 2018). In spite of the care taken to disrupt dynein function at the time of chromosome segregation, the methods used likely targeted all dynein, leaving open the possibility that the lagging chromosomes observed resulted from spindle defects rather than from a failure to move chromosomes along lateral microtubules. Finally, *C. elegans* chromosomes are holocentric, with centromeres and kinetochores dispersed throughout the chromosomes (Cheeseman et al., 2004; Dumont et al., 2010; Hagstrom et al., 2002; Howe et al., 2001). Thus even microtubules that pass laterally by microtubules might make contact with kinetochores.

In support of the Dumont and Desai end-on pushing model, all studies of microtubule dynamics during oocyte meiosis in *C. elegans* agree that the

of any one mechanism may compromise some but not all movements. Indeed, a recent publication suggests that both lateral and end-on microtubule attachments contribute to chromosome positioning and movement during Drosophila oocyte meiosis I (Radford et al., 2015). More extensive genetic studies that simultaneously reduce the function of different combinations of these alternative pathways may clarify how they are integrated to execute meiotic cell division. But ultimately, a thorough understanding of the pleiotropic functions of these proteins will likely require techniques that selectively inactivate individual proteins in specific subcellular regions at specific times. Given recent advances in optogenetics, such approaches may soon become possible.

12. ADVANCING OUR UNDERSTANDING OF OOCYTE MEIOTIC SPINDLE ASSEMBLY: GENETICS AND LIVE CELL IMAGING

Compared to mitosis, our understanding of oocyte meiotic cell division lags considerably. Classical genetic screens in *C. elegans* and *Drosophila* have revealed important players in this fundamental process, and a recent RNA interference screen in mouse oocytes promises to provide further insight (Pfender et al., 2015). While genetic screens and live cell imaging thus far indicate that there are common features among organisms, the molecular pathways that mediate meiotic spindle assembly appear to be numerous and diverse. To advance our understanding, more systematic genetic analyses are needed.

But genetic analysis has its limits. For example, many of the genes required for meiotic spindle assembly have earlier essential requirements.

Gard, D. L. (1992). Microtubule organization during maturation of *Xenopus* oocytes: Assembly and rotation of the meiotic spindle. *Developmental Biology*, *151*, 516–530.

Gomes, J. E., et al. (2013). Microtubule severing by the katanin complex is activated by PPFR-1-dependent MEI-1 dephosphorylation. *The Journal of Cell Biology*, *202*, 431–439.

Gönczy, P. (2012). Towards a molecular architecture of centriole assembly. *Nature Reviews. Molecular Cell Biology*, *13*, 425–435.

Goshima, G., & Kimura, A. (2010). New look inside the spindle: Microtubule-dependent microtubule generation within the spindle. *Current Opinion in Cell Biology*, *22*, 44–49.

Goudeau, M., & Goudeau, H. (1986). The resumption of meiotic maturation of the oocyte of the prawn *Palaemon serratus* is regulated by an increase in extracellular Mg^{2+} during spawning. *Developmental Biology*, *113*, 361–370.

Goudeau, M., & Lachaise, F. (1980). Fine structure and secretion of the capsule enclosing the embryo in a crab (*Carcinus maenas* (L.)). *Tissue and Cell*, *12*, 287–308.

McNally, K., et al. (2014). Katanin maintains meiotic metaphase chromosome alignment and spindle structure in vivo and has multiple effects on microtubules in the Molecular Biology of the Cell, 25, 1037–1049.
McNicoll, F., et al. (2013). Cohesin in gametogenesis. Current Topics in Developmental Biology, 102, 1–34.
Meireles, A. M., et al. (2009). Wac: A new Augmin subunit required for chromosome alignment but not for acentrosomal microtubule assembly in female meiosis. The Journal of Cell Biology, 184, 777–784.
Merdes, A., et al. (2000). Formation of spindle poles by dynein/dynactin-dependent transport of NuMA. The Journal of Cell Biology, 149, 851–862.
Mineyuki, Y. (2007). Plant microtubule studies: Past and present. Journal of Plant Research, 120, 45–51.
Mitchison, T., & Kirschner, M. (1984). Dynamic instability of microtubule growth. Nature, 312, 237–242.
Mitchison, T. J., et al. (2005). Roles of polymerization dynamics, opposed motors, and a tensile element in governing the length of Xenopus extract meiotic spindles. Molecular Biology of the Cell, 16, 3064–3076.
Müller-Reichert, T., et al. (2010). The elegans of spindle assembly. Cellular and Molecular Life Sciences, 67, 2195–2213.
Munro, E., & Bowerman, B. (2009). Cellular symmetry breaking during Caenorhabditis elegans development. Cold Spring Harbor Perspectives in Biology, 1, a003400.
Muscat, C. C., et al. (2015). Kinetochore-independent chromosome segregation driven by lateral microtubule bundles. Elife, 4, e06462.
Musch, A. (2004). Microtubule organization and function in epithelial cells. Traffic, 5, 1–9.
Nakashima, S., & Kato, K. H. (2001). Centriole behavior during meiosis in oocytes of the sea urchin Hemicentrotus pulcherrimus. Development, Growth & Differentiation, 43, 437–445.
Nelson, G. A., et al. (1982). Caenorhabditis elegans spermatozoan locomotion: Amoeboid movement with almost no actin. The Journal of Cell Biology, 92, 121–131.
Nigg, E. A., & Raff, J. W. (2009). Centrioles, centrosomes, and cilia in health and disease. Cell, 139, 663–678.
Oakley, B. R., et al. (2015). gamma-Tubulin complexes in microtubule nucleation and beyond. Molecular Biology of the Cell, 26, 2957–2962.
Ohkura, H. (2015). Meiosis: An overview of key differences from mitosis. Cold Spring Harbor Perspectives in Biology, 7, a015859.
Ori-McKenney, K. M., et al. (2012). Golgi outposts shape dendrite morphology by functioning as sites of acentrosomal microtubule nucleation in neurons. Neuron, 76, 921–930.
O'Rourke, S. M., et al. (2007). Dynein modifiers in C. elegans: Light chains suppress conditional heavy chain mutants. PLoS Genetics, 3, e128.
Paaby, A. B., et al. (2015). Wild worm embryogenesis harbors ubiquitous polygenic modifier variation. Elife, 4, 309178.
Palacios, M. J., et al. (1993). Gamma-tubulin reorganization during mouse fertilization and early development. Journal of Cell Science, 104 (Pt. 2), 383–389.
Petronczki, M., et al. (2003). Un menage a quatre: The molecular biology of chromosome segregation in meiosis. Cell, 112, 423–440.
Petry, S., & Vale, R. D. (2015). Microtubule nucleation at the centrosome and beyond. Nature Cell Biology, 17, 1089–1093.
Petry, S., et al. (2011). Augmin promotes meiotic spindle formation and bipolarity in Xenopus egg extracts. Proceedings of the National Academy of Sciences of the United States of America, 108, 14473–14478.
Pfender, S., et al. (2015). Live imaging RNAi screen reveals genes essential for meiosis in mammalian oocytes. Nature, 524, 239–242.

Phillips, P. C., & Bowerman, B. (2015). Cell biology: Scaling and the emergence of evolutionary cell biology. *Current Biology*, *25*, R223–R225.
Powers, J., et al. (2004). Loss of KLP-19 polar ejection force causes misorientation and missegregation of holocentric chromosomes. *The Journal of Cell Biology*, *166*, 991–1001.
Prosser, S. L., & Pelletier, L. (2015). Centrosome biology: The ins and outs of centrosome assembly. *Current Biology*, *25*, R656–R659.
Radford, S. J., et al. (2012). The chromosomal passenger complex is required for meiotic acentrosomal spindle assembly and chromosome biorientation. *Genetics*, *192*, 417–429.
Radford, S., et al. (2015). Lateral and end-on kinetochore attachments are coordinated to achieve bi-orientation in Drosophila oocytes. *PLOS Genetics*, *11*, e1005605.
Reschen, R. F., et al. (2012). Dgp71WD is required for the assembly of the acentrosomal meiosis I spindle, and is not a general targeting factor for the gamma-TuRC. *Biology Open*, *1*, 422–429.
Saiki, T., & Hamaguchi, Y. (1997). Division of polar bodies induced by their enlargement in the starfish *Asterina pectinifera* egg. *Experimental Cell Research*, *237*, 142–148.
Sampath, S. C., et al. (2004). The chromosomal passenger complex is required for chromatin-induced microtubule stabilization and spindle assembly. *Cell*, *118*, 187–202.
Saunders, A. M., et al. (2007). Kinesin-5 acts as a brake in anaphase spindle elongation. *Current Biology*, *17*, R453–R454.
Sawada, T., & Schatten, G. (1988). Microtubules in ascidian eggs during meiosis, fertilization, and mitosis. *Cell Motility and the Cytoskeleton*, *9*, 219–230.
Sawin, K. E., et al. (1992). Mitotic spindle organization by a plus-end-directed microtubule motor. *Nature*, *359*, 540–543.
Schuh, M., & Ellenberg, J. (2007). Self-organization of MTOCs replaces centrosome function during acentrosomal spindle assembly in live mouse oocytes. *Cell*, *130*, 484–498.
Siegbert, C., et al. (2003). KLCGu(,)-20(,)-2K0rKa53(spindfkinesin0-276((2213(HoR4 uired)-2

Szollosi, D., et al. (1972). Absence of centrioles in the first and second meiotic spindles of mouse oocytes. Journal of Cell Science, 11, 521–541.
Tanaka, N., et al. (2012). Nezha/CAMSAP3 and CAMSAP2 cooperate in epithelial-specific